

User Manual

OriCell™ Balb/c Mouse Embryonic Stem Cells

Catalog No. MUCES-01001



Introduction

Embryonic stem cells (ESCs) are pluripotent stem cells derived from the inner cell mass of a blastocyst. They have the ability to differentiate into ectoderm, endoderm, and mesoderm, giving rise to various cell types. Unlike other stem cells, embryonic stem cells can proliferate indefinitely. Their plasticity and unlimited proliferation make them a major focus in regenerative medicine and tissue engineering research.

OriCell™ Balb/c Mouse Embryonic Stem Cells maintain a normal diploid karyotype after in vitro expansion and culture. They express specific embryonic stem cell markers, can form embryoid bodies (EB) in vitro, and are capable of generating teratomas in vivo. These cells are valuable tools for basic and applied research in fields such as developmental biology, regulatory mechanisms, regenerative medicine, and potential therapeutic applications. Furthermore, genetically modifying these embryonic stem cells and introducing them into the mouse germline is an effective approach to produce genetically modified mice.

OriCell™ Balb/c Mouse Embryonic Stem Cells are derived from the inner cell mass of blastocysts from pregnant Balb/c mice at 3.5 days post-coitum. They are cultured on feeder layers of γ -irradiated mouse embryonic fibroblasts (MEFs) using OriCell™ Complete Medium for Mouse Embryonic Stem Cells (Cat. No.: MUXES-90011).


Note: This product is only provided for further scientific research. It is not intended for diagnostic, therapeutic, clinical, household, or any other applications.

When citing our products in academic journals, please indicate “OriCell™ + Catalog Number, from Cyagen Biosciences (Guangzhou) Inc.”

Product Information

Name	OriCell™ Balb/c Mouse Embryonic Stem Cells
Catalog Number	MUCES-01001
Number of Cells	1×10 ⁶ cells/vial
Passage Number	P20
Storage at	Liquid Nitrogen (-196°C)

The Shape of OriCell™ Balb/c Mouse Embryonic Stem Cells



QC

- Pass the detection of bacteria, fungi, mycoplasma and endotoxins.
- Pass the viability examination. The viable rates is higher than 50%.
- The cell doubling time is less than 72 hours.
- By immunofluorescence detection, it expresses Oct4, SSEA-1 and Nanog ($\geq 90\%$), but does not express SSEA-3 and SSEA-4 ($\leq 5\%$).

Please refer to "COA" for details.

General Handling Principles

1. Ensure that all equipment is kept clean and tidy.
2. Please follow the instructions.
3. Use suitable and reliable consumables and reagents.
4. If using feeder cells for culture, it is recommended to use OriCell™ ICR Mouse Embryonic Fibroblasts (Inactivated) (Cat. No.: MUIEF-01002) .
5. If using serum-free and feeder-free culture conditions, it is recommended to use OriCell™ Serum Free Medium For Mouse Embryonic Stem Cells (Type II, Feeder-free) (Cat. No.: MUXES-90061).
6. Usually the inoculation density of OriCell™ Balb/c Mouse Embryonic Stem Cells (Cat. No.: MUCES-01001) is $(1-2) \times 10^4$ live cells/cm².

Note: The cryopreservation solution of this product contains DMSO, which has potential risks. Please handle it carefully.

Culture Vessel Coated with 0.1% Gelatin

Materials Required

- OriCell™ 0.1% Gelatin Solution (Cat. No.: GLT-11301)

Steps

Note: In order to make the first generation mouse embryonic fibroblasts irradiated by γ -rays more effectively adhere to the culture vessel, the surface of the culture vessel should be coated with gelatin.

1. Add an appropriate amount of 0.1% gelatin to the culture flask to cover the entire bottom surface of the culture flask.

2. Shake the liquid to cover the entire bottom surface of the culture flask.
3. Place the culture flask covered with gelatin on the ultra-clean bench for at least 30 minutes.
4. After 30 minutes, remove the gelatin and wait for the flask to dry before it can be used to inoculate cells.

Note: Culture flasks coated with gelatin can be stored at 4°C for two weeks under sterile conditions also the gelatin is not evaporated to dryness.

Thawing of Cells

Materials Required

- OriCell™ ICR Mouse Embryonic Fibroblasts (Cat. No.: MUIEF-01002)
- OriCell™ Complete Medium For Mouse Embryonic Fibroblasts (Cat. No.: MUXEF-90011)

Steps

Note: If the received cells are thawed within 24 hours, they can be stored in a refrigerator at -80°C. If more than 24 hours, please store them in liquid nitrogen. Please take them out 10 minutes early before thawing and place them at -80°C to allow the liquid nitrogen in the tube to evaporate.

1. Preheat the water bath at 37°C.
2. Warm the complete medium to 37°C.
3. Add more than 5 mL of complete medium to a 15mL centrifuge tube for use.
4. Take the cells out of the -80°C refrigerator, put them in a 37°C water bath and shake them quickly to thaw the cryopreservation solution.

Note: During the thawing process, the cryotube must be shaken to ensure that the solution thaws quickly and evenly.

5. When shaking, please avoid water immersing the pipe cover to cause pollution.

6. When the cryopreservation solution has thawed into ice crystal with a diameter of about 2 mm, stop the water bath. Continue to shake the cryotube until the ice crystal melts thoroughly.
7. Wipe the outer surface of the cryotube with 75% ethanol.
8. Open the cryopreservation tube in the ultraclean bench, use a Pasteur pipette to suck the cell suspension, and transfer it to the prepared centrifuge tube.
9. Wash the cryotube once with 1mL of complete medium to collect residual cells to reduce loss.
10. Centrifuge the cell suspension at 250×g for 4 minutes.
11. Remove the supernatant after centrifugation. Add 1mL of complete medium, gently pipette the cell pellet, blow and mix thoroughly.
12. Inoculate the cells into a culture vessel at a density of 2.5×10^4 viable cells/cm², add a sufficient amount of embryonic fibroblast complete medium, and gently shake the cell culture vessel to distribute the cells evenly.
13. Shake the cells well and incubate them in a CO₂ incubator at saturated humidity, 37°C, 5% CO₂ inside.

Note: Do not move or observe the cells within 2 hours of inoculation. This will seriously affect cell adhesion, resulting in poor shape, cell clumping, and uneven adhesion.

14. On the next day of recovery, observe the cell status, and replace medium with fresh complete medium or passage.

Note:

- If you find lots of floating cells or other abnormal conditions, please investigate the cause in time and contact us.
- MEF cells need to be resuscitated the day before embryonic stem cell resuscitation.
- If the mouse embryonic stem cells will be resuscitated on the day of MEF cell replacement, they can be directly replaced with mouse embryonic stem cell culture medium.
- MEF after resuscitation should be used within 3 days.

Thawing and Culturing of Cells

Materials Required

- OriCell™ Balb/c Mouse Embryonic Stem Cells (Cat. No.: MUCES-01001)
- OriCell™ Complete Medium For Mouse Embryonic Stem Cells (Cat. No.: MUXES-90011)

Steps

Note: If the received cells are thawed within 24 hours, they can be stored in a refrigerator at -80°C. If more than 24 hours, please store them in liquid nitrogen. Please take them out 10 minutes early before thawing and place them at -80°C to allow the liquid nitrogen in the tube to evaporate.

1. Preheat the water bath at 37°C.
2. Warm the complete medium to 37°C.
3. Add more than 5mL of complete medium to a 15mL centrifuge tube for use.
4. Take the cells out of the -80°C refrigerator, put them in a 37°C water bath and shake them quickly to thaw the cryopreservation solution

Note: During the thawing process, the cryotube must be shaken to ensure that the solution thaws quickly and evenly.

5. When shaking, please avoid water immersing the pipe cover to cause pollution.
6. When the cryopreservation solution has thawed into ice crystal with a diameter of about 2 mm, stop the water bath. Continue to shake the cryotube until the ice crystal melts thoroughly.
7. Wipe the outer surface of the cryotube with 75% ethanol.
8. Open the cryopreservation tube in the ultraclean bench, use a Pasteur pipette to suck the cell suspension, and transfer it to the prepared centrifuge tube.
9. Wash the cryotube once with 1mL of complete medium to collect residual cells to reduce loss.
10. Centrifuge the cell suspension at 250×g for 4 minutes.

11. Remove the supernatant after centrifugation. Add 2mL of complete medium, gently pipette the cell pellet, blow and mix thoroughly.
12. Inoculate the cells into a culture vessel at a density of 2.5×10^4 viable cells/cm², add a sufficient amount of embryonic fibroblast complete medium, and gently shake the cell culture vessel to distribute the cells evenly.
13. Shake the cells well and incubate them in a CO₂ incubator at saturated humidity, 37°C, 5% CO₂ inside.

Note: Do not move or observe the cells within 2 hours of inoculation. This will seriously affect cell adhesion, resulting in poor shape, cell clumping, and uneven adhesion.

15. On the next day of recovery, observe the cell status, and replace medium with fresh complete medium or passage.

Note: If you find lots of floating cells or other abnormal conditions, please investigate the cause in time and contact us.

Passaging of Cells

Materials Required

- OriCell™ 0.25% Trypsin-0.04% EDTA Solution (Cat. No.: TEDTA-10001)
- OriCell™ Phosphate-Buffered Saline Solution (1X) (Cat. No.: PBS-10001)
- OriCell™ Complete Medium For Mouse Embryonic Stem Cells (Cat. No.: MUXES-90011)
- OriCell™ Complete Medium For Mouse Embryonic Fibroblasts (Cat. No.: MUXEF-90011)
- Culture Vessel Coated with MEF

Steps

Note:

- 1) Normally, the cells can be cultured for 48 hours before the next passage;

2) Mouse embryonic stem cells must be passaged when the following conditions occur:

- The clones of mouse embryonic stem cells are relatively large, and they have differentiated or are about to differentiate.
- Although mouse embryonic stem cells did not show obvious differentiation, due to the problem of passaging and inoculation density, inter-clonal fusion occurred or was about to occur.

Experimental Preparation:

Prepare feeder cells 1 to 3 days in advance (see Recovery of ICR Mouse Embryonic Fibroblasts).

1. Preheat complete medium, PBS and trypsin to 37°C.
2. Remove the medium in the culture container.
3. Wash the cells twice with PBS (approximately 3mL for T25 flask and 6mL for T75 flask). Please perform relatively slightly and wash thoroughly. Remove the PBS.
4. Add trypsin (approximately 1.5mL for T25 flask and 3mL for T75 flask), spread quickly to ensure full contact with the cells.
5. Observe the cells under a microscope. After about 70%~80% of the cells have shrunk and round, tap the outer wall of the culture vessel to remove the cells from the culture surface.
6. Add Complete Medium For Mouse Embryonic Fibroblasts (approximately 3mL for T25 flask and 6mL for T75 flask) immediately, and then slightly shake the culture container to mix the medium and trypsin quickly to stop the digestion.
7. Use a pipette to suck up the cell suspension, pipetting the bottom surface of the culture container several times, and pipetting down as much as possible of the cells.

Note: The pipetting action should not be violent.

8. Transfer the cell suspension to a centrifuge tube. Wash the container once with PBS (approximately 3mL for T25 flask and 6mL for T75 flask) to collect residual cells.
9. All the collected cell suspensions are centrifuged at 250×g for 4 minutes.

Note: During centrifugation, replace the MEF prepared in advance with Complete Medium For Mouse Embryonic Stem Cells (preheated to 37°C).

10. Remove the supernatant after centrifugation. Add 2mL of Complete Medium For Mouse Embryonic Stem Cells, gently pipette the cell pellet, blow and mix thoroughly.
11. Inoculate the Mouse embryonic stem cells into a suitable culture container at $(1.0\sim 2.0)\times 10^4$ live cells/cm², or adjust the passage ratio according to the actual growth of the cells.
12. Shake the cells well and incubate them in a CO₂ incubator at saturated humidity, 37°C, 5% CO₂ inside.
13. On the next day of passage, observe the cell status. If more floating cells are found, the medium should be changed.
14. After the cell mass grows to a suitable size, it needs to be passaged or frozen.

Cryopreservation of Cells

Materials Required

- OriCell™ NCR Protein-free Cryopreservation Medium For General Use (Cat. No.: NCPF-10001)
- OriCell™ NCR Cryopreservation Medium For General Use (Cat. No.: NCRC-10001)

OriCell™ NCR Cryopreservation Medium clear composition, suitable for cryopreservation of stem cells and primary cells, without affecting cell growth and differentiation potential. No program is required, and the cells can be directly placed in a refrigerator at -80°C, making the operation more convenient and faster.

Steps

Note:

- **The cells need to be replaced with fresh complete medium 24 hours before cryopreservation.**
1. The cells are cryopreserved after growing to appropriate density that can be passaged.
 2. For cell digestion, please refer to OriCell™ Balb/c Mouse Embryonic Stem Cells “Passaging

Steps 2~10”.

3. The cells are uniformly suspended with an appropriate amount of cryopreserved solution, then the supernatant is removed after centrifugation.
4. The cells are divided into cryopreservation tubes based on proportion or quantity.
5. If you choose OriCell™ NCR Cryopreservation Medium, please disperse the cryopreservation tube directly into the refrigerator at -80°C.

Note: During the cryopreservation of cells, especially within 4 hours of the beginning, the refrigerator door should not be opened, which will seriously affect the survival rate of cells.

6. After 8 hours, cells can be transferred to liquid nitrogen for long-term storage.

Note: We suggest that the storage time in the refrigerator at -80°C should not exceed 48 hours.

Cyagen Biosciences (Guangzhou) Inc. reserves all rights to the technical documents of OriCell™ cell culture products. Without the written permission of Cyagen Biosciences (Guangzhou) Inc. any part of this document shall not be adapted or reprinted for other commercial purposes.